

Volume 6, No. 1 (Supplement)
June 2011
ISSN 1823-2140


The National University
with an
INTERNATIONAL REACH



UNIVERSITI
KEBANGSAAN
MALAYSIA
National University of Malaysia

MEDICINE & Health

The Official Journal of The Faculty of Medicine UKM



22nd - 24th July 2011
Equatorial Hotel, Bangi, Selangor,
MALAYSIA

officiated by
Y.B Datuk Rosnah Haji Abdul Rashid Shirlin
Deputy Minister of Health Malaysia

Organised by



SENSITIVITY OF WHOLE BLOOD CLOTTING TIME COMPARED TO ACTIVATED PARTIAL THROMBOPLASTIN TIME

Ika YY, Rahajuningsih DS

Department of Clinical Pathology, Faculty of Medicine, Universitas Indonesia

Background:

Whole blood clotting time (WBCT) was first introduced by Lee and White in 1912 as a method to assess the clotting system. The role of WBCT in the assessment the intrinsic pathway of coagulation system has later been replaced by activated partial thromboplastin time (APTT) due to higher sensitivity and this test can be done by an automatic coagulometer. However, many clinicians still request WBCT as a part of the preoperative screening test. The aim of this study was to evaluate the sensitivity of WBCT compared to APTT in detecting coagulation abnormality.

Materials & Method:

One hundred patients who underwent coagulation tests were enrolled in this cross sectional study. Whole blood clotting time was performed by the Lee and White method, while APTT was performed on automatic coagulometer Sysmex CA 560. The resupormal or abnormal test was based on their reference value. Statistical analysis was done by Fisher exact test.

Results:

Abnormal coagulation was found in 56 and 13 out of 100 subjects by APTT and WBCT, respectively. There was a significant difference between WBCT and APTT in detecting coagulation abnormality with $p < 0.000$. Sixteen out of 100 subjects were haemophiliac, but only 8 patients indicated prolonged WBCT. WBCT started to prolong at 70 seconds of APTT. The sensitivity and specificity of WBCT was only 23.2% and 100%, respectively.

Conclusion:

There was a significant difference between WBCT and APTT in detecting coagulation abnormality. The sensitivity of WBCT in detecting abnormal coagulation was 23.2% compared to APTT. Therefore WBCT is not recommended as a preoperative screening test.

Keywords:

WBCT, APTT, coagulation abnormality, sensitivity.